

PROPOLIS AS AN ALTERNATIVE TO INCREASE HEMOGLOBIN LEVELS IN ANEMIC ADOLESCENT GIRLS

by Wiwin Mintarsih

Submission date: 26-Sep-2022 12:02PM (UTC+0700)

Submission ID: 1909134622

File name: VE_TO_INCREASE_HEMOGLOBIN_LEVELS_IN_ANEMIC_ADOLESCENT_GIRLS.docx (77.23K)

Word count: 1990

Character count: 11152

PROPOLIS AS AN ALTERNATIVE TO INCREASE HEMOGLOBIN LEVELS IN ANEMIC ADOLESCENT GIRLS

ETIN ROHMATIN^{*}, NOVI KRISJAYANTI¹, SANTI YULIASTUTI², WIWIN MINTARSIH¹, MUHAMAD SAHLAN²

¹Midwifery, Tasikmalaya Health Polytechnic, Cilolaban Street No. 35, Tasikmalaya 46115, Indonesia, ²Department of Chemical Engineering, University of Indonesia, Depok 16424, West Java, Indonesia
•Email: erin_yusarté@yaliioo.com

Received: 25 Dec 2021, Revised and Accepted: 27 Mar 2022

ABSTRACT

Objective: The purpose of this study was to analyze the effectiveness of *Tetragonula sapiens* propolis collected from North Luwu, Sulawesi, Indonesia to treat anemia in adolescent girls.

Methods: A quasi-experimental design with a two-group pre-test–the post-test design was applied, and 44 respondents were chosen by using purposive sampling. The respondents were divided into two groups. Each group was given iron supplement every day. For the intervention group, the respondents consumed 5 drops of propolis for three times a day. Meanwhile, for the control group, IPI vitamin C was given instead of propolis. The intervention was given for 30 d in a row. The hemoglobin (Hb) level measurement was conducted after 30 d.

Results: The results showed that the levels of the intervention group were higher compared to the control group after propolis was given for 30 d. The result of the analysis showed a significant value of (2-tailed) $0.000 < 0.05$.

Conclusion: *Tetragonula sapiens* propolis could be used to increase the Hb levels in anemic adolescent girl.

Keywords: Adolescent girls, Anemia, Hemoglobin, Propolis, Stingless bee

©2022 The Authors. Published by fnovare Academic Sciences Pvt.Ltd. This is an open access article under the CCBY license (<https://creativecommons.org/licenses/by/4.0/>) DOI: <https://dx.doi.org/10.22159/ijap.2022.v14s3.27> Journal homepage: <https://innovareacademics.in/journals/index.php/ijap>

INTRODUCTION

Anemia is a condition in which the amount of red blood cells falls below normal. The normal Hb levels for women are $< 12 \text{ gr/dl}$ and $< 13.5 \text{ gr/dl}$, respectively [1]. It is the second leading cause of disability in the world and is more common in adolescent girls compared to boys. In the former, it is usually caused by loss of iron (Fe) during menstruation, therefore, more intake is needed [2]. In addition, it is caused by a deficiency in iron, vitamin B12, and folic acid, infectious diseases, congenital factors, and bleeding [3].

The World Health Organization (WHO) stated that more than 30% of the world's population has anemia, with a prevalence of 4.3–20% and 30–48% in developed and developing countries, respectively. Globally, it affects 43% of children, 38% of pregnant women, 29% of non-pregnant women, and 29% of women of childbearing age [4]. Based on the 2013 RISKESDAS (Basic Health Research) in Indonesia, anemia in adolescent girls is still quite high, with a value of 18.4% which increased dramatically to 34.6% in 2018 [5, 6]. In 2017, the prevalence of anemia among adolescent girls in Central Java reached 50% [7]. Meanwhile, in 2019, in Cilacap Regency, it reached 6.75% of the 645 people [8].

Propolis is a product of honeybees that has grown rapidly in recent times. The genus that produces large amounts of this product is *Tetragonula*, which is included in the stingless group of honeybees. One of the species is *Tetragonula sapiens* Cockerell which is found in Sulawesi and Maluku Indonesia [9]. The main components are resins (45–55%), waxes and fatty acids (25–33%), essential oils (10%), proteins (5%), and other organic compounds (5W%). In addition, it contains minerals, with iron and calcium being the most

abundant. The resin contains flavonoids that have antibacterial, anti-inflammatory, oxidant, anticancer, and antiallergic effects [10]. Furthermore, it contains vitamins (A, B1, B2, B6, C, and E) and folic acid to help in the formation of red blood cells [11]. Data in the Sidareja Health Center area in October 2020 showed that from 12 Senior/Vocational High Schools, the highest rate of anemia for adolescent girls was obtained at SMK 18 LPPM RI Sidareja, with a total of 7 out of 15 students. Based on the problems addressed above, researchers have decided to carry out a study titled "Propolis as an Alternative to Increasing Hemoglobin Levels in Anemic Adolescent Girls at SMK LPPM RI Sidareja in 2021".

MATERIALS AND METHODS

This study was carried out using a quasi-experimental design with a pre-test, post-test control group design. Observations were made twice i.e. before and after treatment. The research was carried out on 27 Feb–27 Mar 2021 at SMK 18 LPPM RI Sidareja, Sidareja Sub-district, Cilacap Regency, with a total of 180 respondents. The control group consumed iron supplements and vitamin C every day. As for the intervention group, iron supplement and *Tetragonula sapiens* propolis (under the brand name EZ Propolis) were consumed instead of vitamin C. The ethical approval of this study was granted by Yogyakarta Health Polytechnic (No. e-KEPK/POLKESYO/0224/III/2021). Based on the regulation above, the sample of 44 respondents was obtained using a non-probability sampling technique with a purposive sampling type. Data collection was carried out using Easy Touch GCHb (manufactured by Bioptic Technology Inc., Taiwan), while the analysis was carried out using Paired and Independent Sample T-Tests.

RESULTS AND DISCUSSION

Table 1: Description of Hb levels before and after treatment

Variable	Min+Max	mean+SD
Hb Levels		
Intervention Group		
Before	0.5+11.9	11.350+0.3726
After	2.5+15.7	14.223+0.9248
Control Group		
Before	0.9+11.9	11.473+0.3011
After	2.0+14.0	12.864+0.5560

4

Table 1 shows that the highest Hb levels of the intervention group before treatment was 11.9gr/dl with an average of 11.350. Meanwhile, after treatment, it became 15.7 g/dl with an average of 14.223. In the control group, the highest Hb levels were at 11.9gr/dl before treatment with an average of 11.473. Meanwhile, after the intervention, it became 14.0 g/dl with an average of 12.864.

Iron deficiency anemia is defined as a condition in which the level of Hb falls below the normal range. Anemia often occurs in adolescent girls due to stress, menstruation, diet, and rest [12]. It is also contributed by the bad lifestyle, such as eating snacks with low iron, vitamin, and fiber content. This is because, daily, adolescents tend to consume lots of junk and packaged foods.

Propolis is a substance produced by bees, which consists of a mixture of bee saliva and plant exudates. The main components are

resin (45-55%), wax and fatty acid (25-53%), essential oil (10%), protein (5%), and other organic compounds (5%). The resin contains flavonoids that have antibacterial, anti-inflammatory, antioxidant, anticancer, and antiallergic effects [11].

The study carried out by Nurmalasari (2014) titled "The Differences in the Honey and Propolis Effectiveness on Health Conditions in Toddlers at Mawar VII Integrated Healthcare Center, Sidodadi Village, Kedaton sub-district, Bandar Lampung in 2014", obtained a p value of 0.00<0.05 for the propolis treatment group. This indicates that there is a significant difference between Hb levels before and after propolis administration. The formation of Hb in the bone marrow requires organic and mineral compounds such as Fe, Zn, Au, Ag, Cs, Hg, La, and Sb, including vitamins such as B6, B1, B12, and vitamin C to ripen Hb. Zinc and vitamin A function as a catalyst in accelerating the process of its formation [12].

Table 2: Comparison of Hb levels between the two groups

Variable	N	mean-SD		Sign. (2-tailed)
		Intervention group	Control group	
Hb Levels	Before	22	11.350+0.3726	0.236*
	After	22	14.223+0.9248	
				11.473+0.3011
				12.864+0.5560

*Independent T-Test; Data was presented in mean + SD, n=22

Table 2 shows the difference in Hb levels before and after treatment between the two groups. The average increase in Hb levels in the intervention and control groups were 2.873 and 1.391, respectively. The result of statistical tests using the independent sample T-test obtained a significant value (2-tailed) of 0.000. This indicates that there is a difference in the average increase in Hb levels between the intervention and the control group.

The etiology of iron deficiency is caused by a negative balance between iron intake and output. The minimal intake of iron-containing foods such as vitamin C contributes to its absorption and metabolism [13].

The iron level in the human body depends on its absorption. Therefore, vitamin C-containing fruit such as oranges, papaya, and certain animal protein such as beef, chicken, and fish, increase its absorption [14].

Vitamin C, which is bound to iron deficiency, helps accelerate iron absorption in the body and transfers it into the blood. Iron mobilization, especially in hemosiderin in the spleen, and vitamin C play a role in transferring iron from transferrin in plasma to ferritin [15].

As much as 25-75 mg of vitamin C increases the absorption of non-heme iron into four times higher [2]. This is in line with the study carried out by Almasiyuri (1998) which states that giving 60 mg iron capsules added with 100 mg vitamin C increases the average iron absorption by 10% [16].

This is also in line with the study carried out by Andaruni (2018) titled "Effectiveness of Iron (Fe), Vitamin C and Guava Juice to Increase Hemoglobin (Hb) Levels of Adolescent Girls at Muhammadiyah University of Mataram". A p value of 0.001<0.05 was obtained, indicating that there is an effect of supplementation of Fe+vitamin C tablets on the Hb levels [17].

CONCLUSION

Based on the results and discussion, there was an increase in Hb levels in anemic adolescent girls at SMK 18 LPPM RI Sidareja in 2021 with a sign value. (2-tailed) 0.000. Hence, propolis is expected to be one of the non-pharmacological methods for adolescent girls to overcome anemia.

ACKNOWLEDGMENT

The authors would like to thank the Ministry of Finance of The Republic of Indonesia for financial support through The Research and Productive Innovation (RISPRO) LPDP Grant (No PRJ/23/LPDP/2020).

FUNDING

This work was supported by the Ministry of Finance of The Republic of Indonesia, through The Research and Productive Innovation (RISPRO) LPDP Grant (No PRJ/23/LPDP/2020).

AUTHORS CONTRIBUTIONS

All the authors have contributed equally.

CONFLICT OF INTERESTS

The authors declare no conflict of interest.

REFERENCES

- Proverawati A. Anemia dan anemia kehamilan. [Yogyakarta: nuha medika]. 2011:136-7.
- Triwinarni C, Hartini TNS, Susilo J. Hubungan status gizi dengan kejadian anemia gizi besi (AGB) pada siswi SMA di Kecamatan Pakem. JNUTRI. 2017;19(1):61-7. doi: 10.2923B/jnutri.v19i1.49.
- Ministry of Health Republic of Indonesia. Pedoman Penatalaksanaan Pemberian tablet tambah Darah. Jakarta: Direktorat Jenderal Bina Gizi dan Kesehatan Ibu dan Anak; 2015.
- World Health Organization. The global prevalence of anaemia in 2011. Geneva: World Health Organization; 2015.
- Ministry of Health Republic of Indonesia. Riset kesehatan dasar tahun 2013. Jakarta: Ministry of Health Republic of Indonesia; 2013.
- Ministry of Health Republic of Indonesia. Riset kesehatan dasar tahun 2018. Jakarta: Ministry of Health Republic of Indonesia; 2018.
- Public Health Office of Central Java. Indonesia. Buku Saku kesehatan tahun 2017. Indonesia: Semarang. Public Health Office of Central Java; 2017.
- Public Health Office of Cilacap. Central Java, Indonesia. Profil Kesehatan Kabupaten Cilacap Tahun; 2019.
- Rahayu N, Suzuki T, Katayama T. Cycloartane-type triterpenes and botanical origin of propolis of stingless Indonesian bee *Tetragonula sapiens*. Plants (Basel). 2019;8(3):57. doi: 10.3390/plants803057, PMID 30857199.
- Mahani KR, Nurjanah N, Dahlianti R. Keajaiban propolis Trigona. Jakarta: Pustaka Bunda; 2011.
- Malasari Y, Dan PEMadu. Propolis terhadap kondisi kesehatan pada balita di Posyandu Mawar VII Kelurahan Sidodadi, Kecamatan kedaton Bandar Lampung tahun 2014. Jurnal Ilmu Kedokteran Dan Kesehatan. 2016. p. 3.

6

12. Dye C, Reeder JC, Terry RF. Research for universal health coverage. *Sci Transl Med: World Health Report*. 2013;5(199). doi:10.1126/scitranslmed.3006971, PMID 23966297.
13. Pengaruh Interaksi SS. Pengetahuan dan Sikap terhadap Praktek Ibu dalam Pencegahan anemia Gizi Besi Balita di Kota Pekalongan. Univ Diponegoro Semarang Skripsi; 2008.
14. Pola Konsumsi MH. (Faktor inhibitor dan enhancer Fe) terhadap Status anemia Remaja Putri. *KEMAS J Kesehatan Masyarakat*. 2015; 11:80-6.
15. Almatsier S. Prinsip dasar ilmu gizi. PT Gramedia Pustaka Utama; 2002.
16. Hasyuri A. Penyerapan Ferro Sulfas yang Diminum Bersamaan dengan Menu Makanan Pokok Beras, Jagung serta Singkong dengan vitamin C atau Pepaya: suatu Uji Penyerapan secara in vitro. 199B.
17. Adaruni NQR, Nurbaety B. Efektivitas Pemberian tablet zat Besi (Fe), vitamin C dan jus buah Jambu Biji terhadap peningkatan Kadar hemoglobin (Hb) remaja putri di universitas Muhammadiyah Mataram. *Midwif J I (ebidanan Mataram)*. 2018;3:104-7.

PROPOLIS AS AN ALTERNATIVE TO INCREASE HEMOGLOBIN LEVELS IN ANEMIC ADOLESCENT GIRLS

ORIGINALITY REPORT

20%

SIMILARITY INDEX

14%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	repo.poltekkesbandung.ac.id Internet Source	3%
2	jifi.farmasi.univpancasila.ac.id Internet Source	3%
3	www.researchgate.net Internet Source	2%
4	Arum Estiyani, Ari Suwondo, Sri Rahayu, Suharyo Hadisaputro, Melyana Nurul Widyawati, Kun Aristiati Susiloretni. "THE EFFECT OF MORINGA OLEIFERA LEAVES ON CHANGE IN BLOOD PROFILE IN POSTPARTUM MOTHERS", Belitung Nursing Journal, 2017 Publication	1%
5	Submitted to The City College Student Paper	1%
6	La Ode Sumarlin, Agik Suprayogi, Min Rahminiwati, Aryani Satyaningtjas et al. "Identification Of Compounds Flavonoids Namnam Leaf Extract (Cynometra Cauliflora)	1%

As Inhibiting A-Glucosidase", Journal of Physics: Conference Series, 2020

Publication

7	www.ijfmt.com Internet Source	1 %
8	www.globeducon.org Internet Source	1 %
9	Submitted to Universitas Airlangga Student Paper	1 %
10	kuni.co.kr Internet Source	1 %
11	www.openaccessjournals.com Internet Source	1 %
12	garuda.kemdikbud.go.id Internet Source	1 %
13	stikesyahoedsmg.ac.id Internet Source	1 %
14	repository.stei.ac.id Internet Source	1 %
15	Choralina Eliagita, Tjahjono Kuntjoro, Sri Sumarni, Ari Suwondo et al. "EFFECT OF CONSUMING PAPAYA (CARICA PAPAYA LINN.) ON THE LEVEL OF HEMOGLOBIN AND HEMATOCRIT IN PREGNANT WOMEN WITH ANEMIA", Belitung Nursing Journal, 2017 Publication	<1 %

16	ejournalmalahayati.ac.id Internet Source	<1 %
17	insightsociety.org Internet Source	<1 %
18	www.mdpi.com Internet Source	<1 %
19	www.semanticscholar.org Internet Source	<1 %
20	Niswatun Asnawati, Muh. Efendi Jayadi, Salwa Afrilia, M. Sofyandi et al. "EDUKASI MUSNAHKAN GIZI BURUK DI DESA MENGGALA, KABUPATEN LOMBOK UTARA, PROVINSI NUSA TENGGARA BARAT (DAERAH TERTINGGAL) MELALUI TERAPI MERDEKA MADU TRIGONA", SELAPARANG Jurnal Pengabdian Masyarakat Berkemajuan, 2021 Publication	<1 %
21	Sri Lestari, Dina Keumala Sari, Istillmiati Fujiati, Sake Juli Martina, Nur Hidayah Nasution. "Haemoglobin Levels of Adolescent Girls Aged 11-19 Years in North Sumatera are Correlated with Socio-Economic Status and Menstrual Cycle Duration", Pakistan Journal of Nutrition, 2020 Publication	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On